

## The Australian Pancreatic Cancer Genome Initiative (APGI) <a href="http://www.pancreaticcancer.net.au">http://www.pancreaticcancer.net.au</a>



## **APGI BioResource – Patient Derived Cell Line Profiles**

## TKCC-05

PDCL Description:	<b>TKCC-05</b> was derived from mouse xenografts initiated from an undifferentiated adenocarcinoma biopsy taken from the primary tumour of a 56-year-old Caucasian male with cancer of the pancreas.
Mutation Status:	<b>TKCC-05</b> has undergone whole genome sequencing and mutation status can be provided upon request.
Organism:	Human (Homo sapiens)
Tissue:	Pancreas, derived from primary tumour site.
Growth Properties:	Adherent
Morphology:	Mesenchymal. The cells are small and can be spindly at low density but form a monolayer at confluence.
Image:	100 μm
Growth Medium:	DMEM/F-12 supplemented with 15mM Hepes, 10ng/mL human recombinant EGF, 0.1IU/mL Insulin, 40ng/mL hydrocortisone, 0.12% glucose and 7.5% Foetal Calf Serum (FCS).
Cell line revival:	Quick-thaw the vial by gently agitating in a 37°C water bath (approx. 2 mins). A centrifugation step to remove the cryoprotectant medium after thawing is necessary for this cell line.



## The Australian Pancreatic Cancer Genome Initiative (APGI) <a href="http://www.pancreaticcancer.net.au">http://www.pancreaticcancer.net.au</a>



Subculturing Procedure:	Medium Renewal: Twice per week.
	Subcultivation Ratio: 1:2-1:6.
	Seeding density: $0.6 \times 10^4$ cells/cm <sup>2</sup> .
	Harvest the cells at 37°C using PBS-EDTA (PBS + 0.25% glucose + 96 $\mu$ M EDTA) for 15 mins followed by 0.05% Trypsin/EDTA for a further 2 minutes. Flasks will require passaging twice weekly at this seeding density. For a more detailed culturing protocol please see additional documents.
	Culture Conditions: Incubate the culture at 37°C with 5% CO <sub>2</sub> .
	Cryoprotectant Medium: 50% FCS + 40% culture medium + 10% DMSO.
Handling Procedure for Frozen Cells:	Upon receipt, frozen vials should be transferred directly into liquid nitrogen storage (if not used immediately). Prolonged storage at -80°C may result in loss of viability.
Availability:	This cell line is available exclusively from the Garvan Institute of Medical Research, Sydney, Australia. If you are interested in obtaining this cell line, please submit an APGI BioResource Application as per the instructions on the APGI website (www.pancreaticcancer.net.au/bioresource-pdcls/).
Depositor:	Dr Marina Pajic (Personalised Cancer Therapeutics Group, Garvan Institute) on behalf of the APGI
Acknowledgements:	Please acknowledge the APGI and The Garvan Institute of Medical Research in all publications and patent applications that make reference to this cell line.
Use Restrictions:	These cells are distributed for research purposes only and are not to be re-distributed to third parties.
Additional Information:	Please refer to <a href="www.pancreaticcancer.net.au/bioresource-pdcls/">www.pancreaticcancer.net.au/bioresource-pdcls/</a> for more detailed subculture, reagent preparation, freezing and reviving TKCC cell line methodology.